



Electromyographic Biofeedback Immediately Increases Quadriceps Corticospinal Excitability

David Florea MS, ATC,* Michelle McLeod MA, ATC,† Phillip Gribble PhD, ATC, FNATA,†
Michael Tevald, PhD, PT,† Brian Pietrosimone, PhD, ATC‡

*United States Naval Academy, Annapolis, MD, †The University of Toledo, Toledo, OH,
‡University of North Carolina at Chapel Hill, Chapel Hill, NC



Introduction

- Quadriceps corticospinal excitability (CSE) pathways are altered following knee injury.¹
- Altered CSE may contribute to chronic neuromuscular dysfunction, such as quadriceps weakness and decreased voluntary activation which may lead to disability² or the development of osteoarthritis.³
- Previous reviews⁴ have identified modalities that influence spinal reflexive pathways; however, there is no accepted modality that specifically targets impaired CSE.
- Electromyographic biofeedback (EMG-BF) is theorized to enhance the quality of muscle contraction by incorporating externally focused visual and/or auditory real-time feedback to improve motor unit recruitment or optimizing motor unit firing rates.^{5,6}
- The ability of EMG-BF to alter CSE remains unknown. Identifying modalities to enhance excitability is important to optimize rehabilitative efforts to decrease disability and preserve long-term joint health.

Purpose

- Determine the immediate effects of EMG-BF during a maximal voluntary isometric contraction (MVIC) on vastus lateralis (VL) CSE and peak torque, compared to a MVIC without EMG-BF (control) in healthy individuals.


Methods-Participants

All volunteers reviewed and signed an informed consent form and were screened for inclusion/exclusion criteria prior to participation. (Table 1)

Table 1. Means ± SD for Participant Demographics (n=15)

Sex	Age (yrs)	Height (cm)	Mass (kg)
Males = 6	21.47	171.45	70.80
Females = 9	± 3.78	± 10.55	± 14.77

Methods-Study Design



Methods-Set Up





Figure 2. Participant set-up for MVIC and AMT/MEP assessments.

- Participants were seated in an isokinetic dynamometer with hips and knees flexed to 90 degrees.
- Ag/Ag-Cl EMG electrodes were affixed over the VL, 10-cm superior to the superior lateral border of the patella.
- A double-coned coil was placed on the scalp over the area corresponding with the motor cortex.

Methods-Intervention

Figure 3. MVIC with EMG-Biofeedback



- Practice trials were performed to obtain maximal root mean square EMG.
- A visual target line was set on the EMG-BF unit 5% above peak EMG.
- Participants were encouraged to exceed the target line during each MVIC.

Methods-Outcome Measures

- Five peak-to-peak motor evoked potentials (MEP) at baseline, collected at 100% AMT during contractions at 5% of MVIC at the beginning of each session.⁷ (Figure 4)
- Five peak-to-peak MEPs collected at 100% AMT during MVICs during the control or EMG-BF conditions.
- Peak torque was recorded during each MVIC at 150 ms prior to TMS during the control or EMG-BF conditions.



Figure 4. Peak-to-Peak Motor Evoked Potential

- Percent change scores $[(\text{post-pre})/\text{pre}] * 100$ were calculated for changes in peak-to-peak MEP amplitudes, that occurred during EMG-BF and control MVICs compared to baseline.
- Dependent t-tests were utilized to determine differences in MEP change scores and peak torque between conditions. Alpha level was set *a priori* at $P \leq 0.05$.

Results

EMG-BF produced statistically significant increases (*) in MEP change scores ($t_{14}=2.926$, Figure 5a.) and increased torque ($t_{14}=2.186$, Figure 5b.) compared to the control condition.

Results

Figure 5a. Differences in MEP Amplitude Change Scores Between Conditions





Figure 5b. Differences in Mean Peak Torque Between Conditions



Discussion

- Increases in VL MEP amplitude during an MVIC when using EMG-BF indicates a gross overall increase CSE.
- External focus of attention, such as that provided by EMG-BF, has been found to produce greater improvements in acquisition and retention of motor skills.⁸
- Increased CSE may be the mechanism responsible for improved strength found following EMG-BF in the past.⁹
- **CLINICAL IMPACT:** EMG-BF used in conjunction with therapeutic exercise may improve outcomes related to muscle strengthening for patients with impaired CSE.

Conclusion

There was increased VL CSE and knee extension torque during MVICs performed with EMG-BF compared to knee extension MVICs performed without EMG-BF in healthy individuals.

References

1. Ericksen HM, Lephley AS, Gribble PA, Pietrosimone BG. Cortical excitability of the quadriceps is decreased in individuals with unilateral anterior cruciate ligament reconstructions. *J Athl Train.* 2011;46(3):5-36.
2. Pietrosimone BG, Lephley AS, Ericksen HM, Gribble PA, Levine J. Quadriceps strength and corticospinal excitability as predictors of disability after anterior cruciate ligament reconstruction. *J Sport Rehabil.* 2013; 22(1):1-6.
3. Pietrosimone BG, McLeod MM, Lephley AS. A theoretical framework for understanding neuromuscular response to lower extremity joint injury. *Sports Health.* 2012; 4(1): 31-5.
4. Harkey MS, Gribble PA, Pietrosimone BG. Disinhibitory interventions and voluntary quadriceps activation: a systematic review. *J Athl Train.* 2014; [epub ahead of print]
5. Lewitt R, Deisinger JA, Wall JR, Ford L, Cassisi JE. EMG feedback-assisted postoperative rehabilitation of minor arthroscopic knee surgeries. *J Sports Med Phys Fitness.* 1995; 35(3):218-223.
6. Croce RV. The effects of EMG biofeedback on strength acquisition: biofeedback and self-regulation. *Biofeedback Self Regul.* 1986; 11(4): 299-310.
7. Luc BA, Lephley AS, Tevald MA, Gribble PA, White DB, Pietrosimone BG. Reliability of corticomotor excitability in leg and thigh musculature at 14 and 28 days. *J Sport Rehabil.* 2013; [epub ahead of print]
8. Pascua L, Wulf G, Lewthwaite R. Additive benefits of external focus and enhanced performance expectancy for motor learning. *J Sports Sci.* [Epub ahead of print].
9. Lephley AS, Gribble PA, Pietrosimone BG. Effects of electromyographic biofeedback on quadriceps strength: a systematic review. *J Strength Cond Res.* 2012; 26(3):873-82.